

COVID-19 Vaccines for Primary School Kids

How does COVID-19 affect kids in primary school?

All children are at risk of getting COVID-19, but the good news is their symptoms are generally mild.



Many children experience a cough, fever, and a runny nose, and only require rest at home, recovering quickly.¹ A very small percentage of children experience a barking cough, prolonged fever, breathing difficulties and abdominal pain, and these children are advised to see their doctor.



Children with underlying health conditions are at higher risk of experiencing severe COVID-19 symptoms. Conditions include asthma, obesity, prematurity, and compromised immune systems.²⁻⁵

How do COVID-19 vaccines work?

In Australia, children receive mRNA COVID-19 vaccines.

An mRNA vaccine is a way to deliver a message to cells in the body.



- For mRNA COVID-19 vaccines, the message delivered to cells is the instructions on how to make just one part of the COVID-19 virus – the “spike protein”.
- Cells then start making this spike protein. Because the message in the mRNA vaccine tells the body to make only the spike protein, and not the entire COVID-19 virus, it can’t make a person sick from COVID-19.
- The immune system recognises the generated spike proteins on the cells as being foreign to the body and starts training itself to fight off the virus. **If a vaccinated person is exposed to COVID-19, the immune system immediately recognises the COVID-19 spike protein and generates a strong and rapid immune response to try and limit the infection.**

Why do children need a COVID-19 vaccine if they’re not getting very sick?

While severe disease and hospitalisation is uncommon in children, it can occur in some cases and vaccinating children helps to protect against this.⁶ Vaccination also benefits the entire family, as well as the community around your child, as it may help to slow the virus spreading to vulnerable people such as elderly grandparents and younger kids who aren’t yet able to be vaccinated. It is important that we let children keep on doing what children do. We have seen the disruption that COVID-19 can have on young lives from a schooling perspective and also on the social and emotional aspects of their lives.^{7,8}



How many doses do kids need? Will they be required to have booster shots?

The **Pfizer** vaccine is recommended for children aged **five to 11 years**. They should receive two doses, given eight weeks apart. The interval can be shortened in special circumstances to a minimum of 3 weeks, for higher risk groups (such as those with medical risk factors for severe illness) in the context of ongoing community transmission. For children aged five to 11 years, each dose is one third of the dose people aged 12 years and older receive. This dosage is based on age, not weight.⁹



Children aged **six to 11 years** can also receive the **Moderna** vaccine. It's recommended they receive two doses, given 8 weeks apart.¹⁰ The interval can also be shortened in special circumstances to a minimum of 4 weeks. Each dose is one half of the dose that people aged 12 years and older receive. This dosage is based on age, not weight.



Children have incredibly robust immune systems that have a fantastic response to the vaccine and, at this stage, it is not recommended that children receive boosters. *However, it is now recommended that severely immunocompromised children aged five or older should receive three initial COVID-19 vaccines (either Pfizer for children aged five years and older, or Moderna for children aged six years and older), with the third dose given 2 to 4 months after the second dose.*^{10,11}

Is vaccinating children safe?



The Pfizer vaccine has undergone rigorous testing in clinical trials and monitoring in the “real world” and has been shown to be **safe** for primary school-aged children.¹²

More primary school-aged children in America have now received a COVID-19 vaccine than the entire population of children this age in Australia, and this has shown us that it is **safe for children**.¹³

Research demonstrates that the **benefit of having a COVID-19 vaccine far outweighs the risks**. It provides vital protection against severe complications and hospitalisation from the virus and significant vaccine-associated side effects are very rare.¹⁴

Some parents have raised concerns about COVID-19 vaccines and fertility. Studies show that there's **no difference in fertility levels** in women or men before and after COVID-19 vaccination.¹⁵⁻¹⁸

Clinical trials of the Moderna vaccine also show that it's as safe in children as it is in adults. At this stage, there is limited “real-world” data available as the vaccine has not yet been used extensively overseas in primary school-aged children.”¹⁰



What side effects might my child have after vaccination?

Many children only experience mild side effects after receiving the Pfizer vaccine such as pain and swelling in the arm, feeling tired, a mild fever, headache and sore muscles and joints.⁹ Side-effects after the Moderna vaccine are also mild, but may be more common than those following Pfizer vaccination.¹⁰

The most common side effects after Moderna vaccination include pain in the arm, swelling of lymph nodes, fever, headache and nausea.¹⁰



There have been rare reports of more serious side effects such as heart inflammation (myocarditis and pericarditis) in children and adults after both the Pfizer and Moderna vaccine.^{19,20}

Heart inflammation is more likely to occur from a COVID-19 infection rather than from the Pfizer vaccine.²¹

If your child complains of a sore chest, shortness of breath or a fast heartbeat within 5 days of their vaccine, please seek medical care.

How effective is the COVID-19 vaccine in this age group?

Research has shown that the **Pfizer vaccine** given to this age group can prevent more than **90% of COVID-19 infections** seven days after receiving the second dose.¹⁴

The Pfizer vaccine is **highly effective against preventing infection and severe complications that require hospitalisation.**¹⁴



*The clinical trial of the **Moderna vaccine** showed that it creates a similar immune response in children aged six to 11 years when compared to young adults.¹⁰ Though there is limited “real world” data at this stage on how effective the Moderna vaccine is for children, it is likely that Moderna will be very effective at reducing the likelihood of severe COVID-19 in children, including against Omicron.¹⁰*

Where can my child get vaccinated?



In WA, children can be vaccinated at:

- GP clinics,
- state-run clinics,
- some pharmacies and
- Aboriginal Medical Services.

Visit **Roll Up WA** for more information and to make a booking.



For more information on
COVID-19 in kids visit
tacklingcovid19.org.au

References

1. Molteni E, Sudre CH, Canas LS, et al. Illness duration and symptom profile in symptomatic UK school-aged children tested for SARS-CoV-2. *The Lancet Child & Adolescent Health*. 2021;5(10):708-718 Williams P, Koirala A, Saravanos G, et al. COVID-19 in children in NSW, Australia, during the 2021 Delta outbreak: severity and disease spectrum. medRxiv. 2022.
2. Williams P, Koirala A, Saravanos G, et al. COVID-19 in children in NSW, Australia, during the 2021 Delta outbreak: severity and disease spectrum. medRxiv. 2022
3. Hobbs CV, Woodworth K, Young CC, et al. Frequency, characteristics and complications of COVID-19 in hospitalized Infants. *The Pediatric Infectious Disease Journal*. 2021.
4. Mukkada S, Bhakta N, Chantada GL, et al. Global characteristics and outcomes of SARS-CoV-2 infection in children and adolescents with cancer (GRCCC): a cohort study. *The Lancet Oncology*. 2021;22(10):1416-1426.
5. Haeusler GM, Ammann RA, Carlesse F, et al. SARS-CoV-2 in children with cancer or after haematopoietic stem cell transplant: An analysis of 131 patients. *European Journal of Cancer*. 2021;159:78-86
6. NSW Health. COVID-19 monitor: COVID-19 cases, variants, vaccines, hospitalisations and deaths 2022 [cited 2022 Feb]. Available from: https://aci.health.nsw.gov.au/__data/assets/pdf_file/0006/698388/20220104-COVID-19-Monitor.pdf?fbclid=IwAR0MP3YtaypZy3QNYyQHMEPgC-p02Bkz69byB3PMk-t5NtcUKsoi2W1zX_4
7. WA Department of Education, WA Department of Health, Telethon Kids Institute. DETECT schools study: understanding the impact of COVID-19 in Western Australian schools 2021 [cited 2022 Jan]. Available from: <https://www.telethonkids.org.au/globalassets/media/documents/projects/detect-schools-study-final-report-and-appendices.pdf>
8. Racine N, McArthur BA, Cooke JE, et al. Global Prevalence of Depressive and Anxiety Symptoms in Children and Adolescents During COVID-19: A Meta-analysis. *JAMA Pediatrics*. 2021;175(11):1142-1150.
9. Australian Technical Advisory Group on Immunisation (ATAGI). ATAGI recommendations on the use of the paediatric Pfizer COVID-19 vaccine in children aged 5 to 11 years in Australia: Australian Government; 2021 [cited 2022 Feb]. Available from: <https://www.health.gov.au/resources/publications/atagi-recommendations-on-pfizer-covid-19-vaccine-use-in-children-aged-5-to-11-years>
10. Australian Technical Advisory Group on Immunisation (ATAGI). ATAGI recommendations on the use of Spikevax (Moderna) COVID-19 vaccine in children aged 6 to 11 years: Australian Government; 2022 [cited 2022 Feb]. Available from: <https://www.health.gov.au/news/atagi-recommendations-on-the-use-of-spikevax-moderna-covid-19-vaccine-in-children-aged-6-to-11-years>
11. Australian Technical Advisory Group on Immunisation (ATAGI). ATAGI recommendations on the use of a third primary dose of COVID-19 vaccine in individuals who are severely immunocompromised: Australian Government; 2022 [cited 2022 Feb]. Available from: <https://www.health.gov.au/resources/publications/atagi-recommendations-on-the-use-of-a-third-primary-dose-of-covid-19-vaccine-in-individuals-who-are-severely-immunocompromised>
12. AusVaxSafety. COVID-19 vaccines 2021 [cited 2022 Feb]. Available from: <https://ausvaxsafety.org.au/safety-data/covid-19-vaccines>
13. Hause A, Baggs J, Marquez P, et al. COVID-19 Vaccine Safety in Children Aged 5–11 Years—United States, November 3–December 19, 2021. *MMWR Morbidity and mortality weekly report*. 2021;70:1755-1760.
14. Walter EB, Talaat KR, Sabharwal C, et al. Evaluation of the BNT162b2 Covid-19 vaccine in children 5 to 11 years of age. *New England Journal of Medicine*. 2021;386:35-46
15. Bentov Y, Beharier O, Moav-Zafir A, et al. Ovarian follicular function is not altered by SARS–CoV-2 infection or BNT162b2 mRNA COVID-19 vaccination. *Human Reproduction*. 2021;36(9):2506-2513
16. Orvieto R, Noach-Hirsh M, Segev-Zahav A, et al. Does mRNA SARS-CoV-2 vaccine influence patients' performance during IVF-ET cycle? *Reproductive Biology and Endocrinology*. 2021;19(1):69
17. Morris RS. SARS-CoV-2 spike protein seropositivity from vaccination or infection does not cause sterility. *F&S Reports*. 2021
18. Gonzalez DC, Nassau DE, Khodamoradi K, et al. Sperm Parameters Before and After COVID-19 mRNA Vaccination. *JAMA*. 2021;326(3):273-274
19. Therapeutic Goods Administration. COVID-19 vaccine weekly safety report - 06-01-2022: Australian Government; 2022 [cited 2022 Jan]. Available from: <https://www.tga.gov.au/periodic/covid-19-vaccine-weekly-safety-report-06-01-2022>
20. Australian Technical Advisory Group on Immunisation (ATAGI), Cardiac Society of Australia and New Zealand (CSANZ), Royal Australian College of General Practitioners (RACGP), et al. Guidance on myocarditis and pericarditis after mRNA COVID-19 vaccines: Australian Government; 2021 [cited 2022 Feb]. Available from: https://www.health.gov.au/sites/default/files/documents/2021/11/covid-19-vaccination-guidance-on-myocarditis-and-pericarditis-after-mrna-covid-19-vaccines_1.pdf
21. Singer ME, Taub IB, Kaelber DC. Risk of Myocarditis from COVID-19 Infection in People Under Age 20: A Population-Based Analysis. medRxiv. 2021
22. Attwell K, Carlson SJ, Tchilingirian J, Harper T, McKenzie L, Roberts L, Rizzi M, Westphal D, Effler P, Hughes C, Swift V, Blyth CC. Coronavax: Preparing Community and Government for COVID-19 Vaccination: a Research Protocol for a mixed-methods social research project. *BMJ Open*. 2021;11(6):e049356